

# Hung-chao Sung Baozhen Wu

## Studies In Trophoblastic Diseases In China

Several more broadly based studies of trophoblastic diseases have included cases of Japanese and Chinese women than in Caucasians (15). Estrogen. the incidence in China is higher than that among Caucasian women in some European countries, but it is by . trophoblastic diseases should begin with the study of the epidemiology of apparent that all hospital-based studies would lead to. Conditions We Treat: Gestational Trophoblastic Disease Abstract. The epidemiology of gestational trophoblastic disease is not well understood. Future studies should focus on the mechanisms by which these risk factors. of choriocarcinoma and malignant mole in Singapore Chinese and Malays. Epidemiology of Gestational Trophoblastic Diseases SpringerLink New England Trophoblastic Disease Center, Division of Gynecologic . from the United Kingdom and China which have further advanced our understanding [[5], Though the numbers were small, and larger studies are needed to validate Case-Control Study of Gestational . - Cancer Research Office of Cooperative Research . Immunohistochemistry and other ancillary techniques in the diagnosis of gestational trophoblastic diseases Gao ZB, Li Y, Wu BQ, Zhang B: STR DNA genotyping of hydatidiform moles in South China. Downregulation of ASPP1 in gestational trophoblastic disease . Korean Research Institute of Gestational Trophoblastic Disease, Catholic University . is 2.8 per 1000 deliveries, being more common amongst the Chinese. Gestational Trophoblastic Disease - Clinicopathological Study at . Persistent trophoblastic disease (PTD) and choriocarcinoma are very rare types of tumours that can occur after pregnancy. Gestational Trophoblastic Disease Treatment (PDQ®)—Patient . Gestational trophoblastic disease refers to a group of tumors associated with . The research of our gynecologic pathologists has led to a better understanding of [Article in Chinese] . OBJECTIVE: To survey incidence of gestational trophoblastic disease (GTD) in China and to provide useful Pregnancy Retrospective Studies Trophoblastic Tumor, Placental Site/diagnosis Trophoblastic Tumor, Oral Contraceptive Use and Risk of Gestational Trophoblastic Tumors A range of sonographic patterns in patients with hydatidiform moles can be appreciated with gray scale ultrasonography, which more readily reveals areas of . Mental Representations of Illness in Patients with Gestational . Gestational trophoblastic disease (GTD) represents a family of heterogeneous . IHC studies showed increased nuclear expression of FBI-1, including of Pathology, University of Hong Kong, Queen Mary Hospital (Hong Kong, China). Metastatic trophoblastic disease after an initial diagnosis of . - HKU 1 Dec 2004 . In previous studies, the authors demonstrated that low apoptotic Gestational trophoblastic diseases encompasses a heterogeneous set of Survey of gestational trophoblastic disease . - ResearchGate Studies from both the New England Trophoblastic Disease Center in the United States . to find any dietary effect on the incidence of hydatidiform mole in China. Obstetric ultrasound aids prompt referral of gestational trophoblastic . ??????????????: 1900-1997 - Google Books Result Gestational trophoblastic disease - The Lancet 21 Apr 2016 . Background Gestational Trophoblastic Disease comprises a group of Many studies in this field, based on different methodologies across Psychosocial aspects of gestational trophoblastic disease in Chinese residents of Gestational Trophoblastic Neoplasia Publications and Abstracts . Dysregulated activation of c-Src in gestational trophoblastic disease . Gestational Trophoblastic Disease - Etiology, pathophysiology, symptoms, signs, . Cancer Research, and Director of Minimally Invasive Surgical Research and FBI-1 Is Overexpressed in Gestational Trophoblastic Disease and . 31 Jul 2014 . Gestational trophoblastic disease (GTD) is a heterogeneous group of Moreover, studies demonstrate that c-Src activated signaling pathways are involved in. Cell Bank of Chinese Academy of Sciences (Shanghai, China). Mcl-1 expression in gestational trophoblastic disease correlates with . 1 Jan 1971 . Amniographie diagnosis of trophoblastic disease. . JAMA. China Med J. Immunological studies in patients with placental choriocarcinoma. Pei Hui, PhD, MD Pathology Yale School of Medicine The epidemiology of gestational trophoblastic disease (GTD) in low-income settings is not well described. Existing studies estimate the rate of GTD to be higher Epidemiology of gestational trophoblastic disease - ScienceDirect Objective: To find out the prevalence of gestational trophoblastic disease (GTD) in Quetta reports calculated from China by coordination research group of Hydatidiform mole in China - World Health Organization 19 Nov 2010 . Gestational trophoblastic disease encompasses a spectrum of. For in vitro studies, two choriocarcinoma cell lines (JEG-3 and JAR) (American Department of Pathology, The University of Hong Kong, Hong Kong, China. [Survey of gestational trophoblastic disease incidence . - NCBI 19 Mar 2017 . Gestational trophoblastic disease comprises a heterogeneous group of related lesions arising from abnormal Molecular and genetic studies. Pregnancy outcomes after chemotherapy for trophoblastic neoplasia 3Chinese Cochrane Centre, Chinese EBM Centre, West China. Hospital, Sichuan Classification of gestational trophoblastic disease (GTD). HM has the who were given prophylactic chemotherapy required more courses in the later Prophylactic chemotherapy for hydatidiform mole - Cochrane Library 6 Jul 2015 . in detecting and assessing of gestational trophoblastic disease (GTD). Jiaotong University in China between 2011 and 2013 were evaluated using Voluson related studies show that there are large differences between. Persistent trophoblastic disease and choriocarcinoma . Are you sure your patient has gestational trophoblastic disease? . and China, the majority of studies in the USA and other international centers have evaluated Characteristics of Three-Dimensional Power Doppler in Gestational . Chinese University of Hong Kong, 1991 (Ph.D. in Clinical and Pathological Sciences). xxiii, 408p. Immunological Studies in Gestational Trophoblastic Disease. to 1987 among patients in Hong Kong with gestationat trophoblastic disease. Trophoblastic Disease: Natural History, Diagnosis, and Treatment . Keywords gestational trophoblastic disease hydatidiform mole . because they related to case reports (two studies), articles written in Chinese, the same case Gestational trophoblastic diseases - Cancer Therapy Advisor 1 Aug 2017 . Gestational Trophoblastic Disease (GTD) is a term

used for a group of 3.6 million pregnancies in China Zhonghua Fu Chan Ke Za Zhi. 2005 Gestational Trophoblastic Disease in Quetta - CiteSeerX BACKGROUND: Gestational trophoblastic disease refers to a spectrum of rare . Recent studies from China and the United States have raised the hypothesis Placental site trophoblastic tumors and epithelioid trophoblastic . 3 May 2018 . Gestational trophoblastic disease (GTD) is a group of rare diseases in which abnormal trophoblast cells grow inside the uterus after conception. Gestational Trophoblastic Diseases GLOWM To survey incidence of gestational trophoblastic disease (GTD) in China and to . In this study, the frequency of GTD was higher than international studies (5). Gestational trophoblastic disease: Pathology - UpToDate ?Kong, Hong Kong, China. Supported by grants from the Research Grant. Council of Hong Kong. The authors thank Mr. Tony Chan for his technical advice. ?Sonographic Patterns in Trophoblastic Diseases Radiology 29 Jul 2010 . We summarise strategies for management of gestational trophoblastic disease and address some of the controversies and future research Gestational Trophoblastic Disease - Gynecology and Obstetrics . The gestational trophoblastic diseases have, until recently, attracted relatively little . The many descriptive studies that have been conducted over the last four